

175mm deep

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE USING
FOR ANIMAL TREATMENT ONLY

LIFT
HERE



Pour-On Lousicide For Sheep
35 g/L IMIDACLOPRID

For the control of susceptible body lice in *Bovicola Ovis* on shorn sheep off shears and unshorn lambs up to 2 months of age. This product can be expected to protect sheep against re-infestation with lice for 4 weeks after treatment.




L2852 0610

20 Litres APVMA Approval No 62598/20L/0510 

200mm wide

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE USING
FOR ANIMAL TREATMENT ONLY


LIFT
HERE





AVENGETM

Pour-On Lousicide For Sheep
35 g/L IMIDACLOPRID

For the control of susceptible body lice in *Bovicola Ovis* on shorn sheep off shears and unshorn lambs up to 2 months of age. This product can be expected to protect sheep against re-infestation with lice for 4 weeks after treatment.



L2852 0610

 **20 Litres** APVMA Approval No 62598/20L/0510 

200mm wide

175mm deep

Job: 175911 **YOUR PROOF**
Name: 02 Avenue 20L Top
Client: Bayer Australia
Date: 1/06/2010








CHECK A PANTONE BOOK FOR TRUE COLOUR REPRESENTATION

OK to proceed
 Further proof required

sign: _____ date: _____

No responsibility is taken for errors not noted on this proof
EXAMINE THIS ART VERY CAREFULLY AND RETURN THE SIGNED PROOF TO US FOR THE JOB TO PROCEED

PHILSTIC Philstic Labels Limited
Email: prepress@philstic.co.nz
Phone: +64 9 970 1430
Fax: +64 9 630 0957

-  Dieline
-  Process Blue
-  PMS 2935
-  PMS 368
-  PMS 200
-  Black
-  White

AVENGE™ 35g/L POUR-ON LOUSICIDE FOR SHEEP

AVENGE™ POUR ON is for the control of neonicotinoid susceptible sheep body lice *Bovicola ovis* in sheep off-shears (within 24 hours of shearing) and unshorn lambs up to 2 months of age.

DIRECTIONS FOR USE

This product can be used on all adult sheep up to 24 hours off-shears and on unshorn lambs up to 2 months of age.

Apply the entire dose to sheep under 30kg as a single broadband stripe from poll to tail butt.

For sheep over 30kg, apply half the dose as a broadband stripe either side of the midline from the mid neck to the butt of the tail. The stripes should be no further than 10 cm apart. Refer to the diagrams below. A representative sample of animals should be weighed before treatment. Dose the mob to the heaviest animal by liveweight in each group (ewes, wethers, rams, lambs). Do not underdose. Where there is a large variation in size within (e.g lambs) the group, dose rate should be based on the label directions for each weight range. Drafting into two or more lines may be appropriate, to avoid excessive overdosing.

Bodyweight (kg)	Dose (mL)	Method	Apply in the pattern below
Below 6kg	8	Single stripe	
6.0 - 8.0	12		
8.1 - 10.0	15		
10.1 - 12.5	20		
12.6 - 15	25		
15.1 - 20	30	Double stripe	
20.1 - 30	40		
30.1 - 55	60		
55.1 - 80	80		
80.1 - 100	90		

Add 10 mL for each 10 kg above 100 kg

WITHOLDING PERIODS

MEAT: DO NOT USE less than 21 days before slaughter for human consumption.

MILK: DO NOT USE on female sheep which are producing, or may in the future produce, milk or milk products for human consumption.

WOOL: DO NOT USE less than 6 months before shearing or fibre collection.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

EXPORT SLAUGHTER INTERVAL (ESI): DO NOT slaughter for export less than 63 days after treatment. The ESI on this label was correct at the time of label approval. Before using this product, confirm the current ESI from the Bayer Website www.bayeranimal.com.au or via the APVMA website at www.apvma.gov.au/residues/ESI.shtml or by calling Bayer Australia Ltd on 1800 678 368.

SAFETY DIRECTIONS

May irritate the eyes. Avoid contact with eyes. When using the product wear face shield or goggles. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water.

Rehandling Interval: Do not rehandle sheep after treatment until dry.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia, 131126. Additional information is listed in the Material Safety Data Sheet.

AVENGE™ is a pending trademark of Bayer AG, Leverkusen, Germany.

DRUM TAP ENCLOSED
Fit to internal thread after removing small drum cap



Cut along dotted line.

CUSTOMER INFORMATION LINE
1800 678 368
 TOLL FREE from anywhere in Australia
 9:00am to 4pm (Eastern Time)
 Monday to Friday
www.bayeranimal.com.au

Storage: Store below 30°C (Room temperature) Store in original container, tightly closed in a safe place.

See container stamping for batch and expiry date.
 APVMA Approval No 62598/20L/0510

© Bayer Australia Ltd 2008
 Manufactured in New Zealand

Bayer Australia Ltd. ACN 000 138 714.
 875 Pacific Highway, PYMBLE, NSW, 2073



L2853 0610

Job: 175911 **YOUR PROOF**
Name: 02 Avenue 20L Back
Client: Bayer Australia
Date: 1/06/2010

CHECK A PANTONE BOOK FOR TRUE COLOUR REPRESENTATION

OK to proceed
 Further proof required

sign: _____ date: _____

No responsibility is taken for errors not noted on this proof
EXAMINE THIS ART VERY CAREFULLY AND RETURN THE SIGNED PROOF TO US FOR THE JOB TO PROCEED

PHILSTIC Philstic Labels Limited
 Email: prepress@philstic.co.nz
 Phone: +64 9 970 1430
 Fax: +64 9 630 0957

- Dieline
- PMS 2935
- Black

175mm deep

200mm wide

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE USING
FOR ANIMAL TREATMENT ONLY

AVENGE™ Pour-On Lousicide For Sheep
35g/L IMIDACLOPRID

General Instructions

Apply product only with the Bayer 6-stripe applicator. Treated sheep must not be mixed with lice free sheep until 6 weeks after treatment. Live lice on ewes treated within 6 weeks of lambing can infect lambs from these ewes.

Sheep must be treated within 24 hours of shearing. Unshorn lambs can be treated up to 2 months of age. Efficacy is not affected by moderate rainfall either before or after treatment (50 mm rain 1 hour prior to treatment and/or 12.5 mm 2 hours after treatment and/or 25 mm 24 hours after treatment). Treated sheep must not be mixed with lousy sheep. Do not use on sheep which have not been cleanly shorn for any reason eg., mycotic dermatitis.

Management practices which promote good lice control such as complete musters, accurate treatment of every animal according to label, ensuring all animals can be treated at the same time, no split shearings, stock-proof fencing to prevent external introductions, quarantine and lice treatment of all newly purchased stock are essential for effective lice control.

Treated sheep could potentially develop backline dermatitis following prolonged exposure to extreme heat and sunlight without access to shade. Do not leave container, hose or applicator in the sun for long periods of time.

AVENGE™ POUR ON may be used in conjunction with worm control drenches and at the same time as vaccination.

AVENGE™ POUR ON contains a blue dye to identify treated sheep. This dye is scourable from the wool. The product can be detrimental to certain plastics, paints and surface finishes. Avoid contact with such surfaces.

AVENGE™ POUR ON is safe to use on breeding rams.

WITHHOLDING PERIODS

MEAT: DO NOT USE less than 21 days before slaughter for human consumption.

MILK: DO NOT USE on female sheep which are producing, or may in the future produce, milk or milk products for human consumption.

WOOL: DO NOT USE less than 6 months before shearing or fibre collection.

EXPORT SLAUGHTER INTERVAL (ESI): DO NOT slaughter for export less than 63 days after treatment.

The ESI on this label was correct at the time of label approval. Before using this product, confirm the current ESI from the Bayer Website www.bayeranimal.com.au or via the APVMA website at www.apvma.gov.au/residues/ESI.shtml or by calling Bayer Australia Ltd on 1800 678 368.

SAFETY DIRECTIONS

May irritate the eyes. Avoid contact with eyes. When using the product wear face shield or goggles. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands arms and face thoroughly with soap and water.

First Aid

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia, 131126. Additional Information is listed in the Material Safety Data Sheet.

Rehandling Interval: Do not rehandle sheep after treatment until dry.

Protection of Wildlife, Fish, Crustaceans and the Environment

Imidacloprid is highly toxic to aquatic species. Do not contaminate dams, ponds, rivers, waterways or drains with pesticide or used containers.

Disposal of Containers

For disposal containers must be triple rinsed with diesel, and dispose of rinsate in a disposal pit away from desirable plants and their roots, and watercourses. On site disposal of undiluted chemicals is unacceptable. If not recycling break, crush or puncture and return empty packaging for appropriate disposal in a local authority waste management facility. Do not burn empty containers.

Limitation of Liability

Bayer Australia Limited (BAL) warrants that this product is of merchantable quality and fit for its intended purpose indicated on this label. Failure to use the product strictly as directed may be illegal, prove dangerous or render the product ineffective. Notwithstanding anything else contained on this label BAL's liability is limited to replacing this product or supplying equivalent product or paying the cost of replacing this product, at its discretion.

Bayer Australia Ltd.
875 Pacific Highway,
PYMBLE, NSW, 2073
ACN 000 138 714



Storage: Store below 30°C (Room temperature) Store in original container, tightly closed in a safe place.

See container stamping for batch and expiry date.

APVMA Approval No 62598/20L/0510

© Bayer Australia Ltd 2008

Manufactured in New Zealand

L2852 0610

173mm deep

175mm wide

175mm wide

Job: 175911	YOUR PROOF
Name: 02 Avenge 20L Book	
Client: Bayer Australia	
Date: 1/06/2010	
CHECK A PANTONE BOOK FOR TRUE COLOUR REPRESENTATION	
<input type="checkbox"/> OK to proceed	
<input type="checkbox"/> Further proof required	
sign: _____	date: _____
No responsibility is taken for errors not noted on this proof	
EXAMINE THIS ART VERY CAREFULLY AND RETURN THE SIGNED PROOF TO US FOR THE JOB TO PROCEED	



Philstic Labels Limited
Email: prepress@philstic.co.nz
Phone: +64 9 970 1430
Fax: +64 9 630 0957



Dieline



Black