

AVENGE™

Break resistance.

Tips to achieve and maintain a lice-free flock using AVENGE.

AVENGE is a knockdown product, so the prevention of reinfestation is important.

Management tips.

- Start with a complete muster. Lice are only acquired from other sheep, so a second muster is recommended in rough country.
- Clean shearing is important. If cotted fleeces, grass seeds, lumpy wool or flystrike prevent clean shearing, lice control by any method will be poor.
- Prevent reinfestation. Avoid split shearing. Maintain boundary fences in good repair. Shear, treat and quarantine any introduced sheep whose lice status is not known.
- Pregnant ewes may be treated up to 6 weeks before lambing. Lice from ewes can infest young lambs.
- Treated sheep should not be mixed with untreated lice-free sheep for at least six weeks.
- AVENGE is a knockdown product, so the prevention of reinfestation is important. Check sheep for signs of re-infestation at five to six months. If reinfestation has occurred, your Bayer Area Manager can assist you with a control strategy.

Apply AVENGE to sheep within one day of shearing.

Application tips.

- Apply AVENGE to sheep within one day of shearing.
- Before applying AVENGE, shake the container and check the dose accuracy of the applicator gun.
- Dose to the heaviest bodyweight in the mob. Check by weighing, don't guess.
- Apply AVENGE in continuous broad-band stripes along the backline.
- Check dye marks on all sheep at the end of each race load. Re-treat any animals in doubt.
- AVENGE is rainfast. However, it is Good Veterinary Practice to treat dry sheep and avoid treating in heavy rainfall periods if possible.
- Clean applicator gun thoroughly following use.
- Store AVENGE below 30°C (room temperature) in original container, tightly closed in a safe place.

